

Directions of APIFLUME Strips

Veterinary OTC Medicine for External Use


【Generic Name】 Flumethrin Strip

【Proprietary Name】 APIFLUME

【Composition】 Flumethrin

【Molecular Formula】 $C_{28}H_{22}Cl_2FNO_3$

【Structural Formula】



【Description】 It occurs as a yellow to orange strip,slightly foul odor.

【Pharmacological Actions】

Flumethrin acts on the nervous system of pests, disrupting the function of the nervous system through the action of sodium ion channels. This product is a pyrethroid pesticide. It is mainly used for the control of ectoparasites of bees and livestock, and to inhibit the activity of adult egg laying and egg hatching.

【Drug Interaction】 Do not use it with other pesticides or alkaline drugs because it may be invalid.

【Indication】 Miticide,which is used to kill and control varroa mites.

【Dosage & Administration】

1. Hanging Method: Hang the strip between two comb frames just over the beehive entrance or between two comb frames diagonally.Apply 1/3~1/2 strip to beehive with less than 4 comb frames;Apply 1/2~2/3 strip to beehive with 4~8 comb frames;Apply 2/3~1 strip to beehive with more than 8 comb frames;Apply the strips one more time after 5~7 days.Applying medicine 3~4 times is a course of treatment.
2. Trial is required. Choose 3 or 5 bee colonies for trial before applying to the whole colonies. Make a specific medicine using plan after observing 24~48 hours of the trial. It's better to conduct the trial in the morning of sunny day and adjust dose according to actual situation.Increase or reduce dose or medicine applying time on the condition that there are step hives,less comb frames or severe varroa mites disease.

【Adverse Reactions】

This product has no obvious adverse reactions to healthy bee colonies; It may cause crawling bees for bee colonies with poor natural resistance (such as Megachile), or unhealthy bees, sensitive bee colonies, and bees at different developmental stages with differences in drug-resistance. After taking medicine, observe for several hours and take some necessary actions.

【Precautions】

1. Please air-dry the strips in the shade for 24~48 hours if you find the strips are wet or sticky to each other. Strips can be come into use when there's no stickiness to fingers, otherwise it may cause harm to bees. Please do not place it under the Sun, otherwise the efficacy will be greatly decreasing.
2. Do not use this product when temperature is above 35℃ or below 15℃. The product is not recommended to be applied to the situation such as migratory beekeeping, the beginning of hiving bee colonies, sugar deficiency for bees or other abnormal situation .
3. The said product can not be applied to bee colonies that have been fed with alkaline pollen such as China camellia pollen or other alkaline medicines and products recently or in the period of camellia collection.
4. The strips should not be placed continuously in beehive for more than 6 weeks.
5. When one decides to hang strips, please wear protective gloves and mask. Wash hands after use.
6. Disused strips should not be reused and should be disposed in a centralized place.
7. The strips are prohibited from contact with edible honey and should not be stored together with pesticides or other chemicals.
8. Store the said product somewhere away from food and children's reach and only unpack the packet when it is near to use.
9. Do not use this product during honey flow period.

【Withdrawal Time】 It is not required to be determined.

【Specification】 3.6 mg/strip.

【Shelf Life】 4 years.

【Storage】 Keep it sealed and away from light

【Manufacturer】 Sichuan Lingman Biotechnology Co.,Ltd.

【Address】 No.10, Yugui Road, New District, Economic Development Zone, Meishan City, Sichuan Province, China

【Sales Company】 Chengdu Flybee Trading Co.,Ltd.

【Website】 www.fly-bee.com

【Email】 sales_02@fly-bee.com

Introduction to Graphical Method

APIBUZZ BEE MEDICINE **Anti-Varroa Mite Strip**



FEATURES OF APIBUZZ BEE MEDICINES

Harmless to Bees

Better Stability

High Efficacy

Convenient to Use

No Residual



BEFORE USE




AFTER USE



Product Detail

Product Name ----- Apiflume Flumethrin 20-Strip
 Generic Name ----- Flumethrin Strip
 Brand ----- Apibuzz
 Packing Size ----- -20 Strips/Packet
 Strip Size ----- -21cm x 3.5cm
 Active Content ----- Flumethrin
 Efficacy ----- The said product is for bee mite prevention and treatment. It has a strong killing effect on .bee varroa mites with strong drug resistance
 Expiry Time ----- 4 Years





Dosage & Administration

HANGING METHOD

Hang the strip between two comb frames just over the beehive entrance or between two comb frames diagonally. Apply $\frac{1}{3}$ ~ $\frac{1}{2}$ strip to beehive with less than 4 comb frames; Apply $\frac{1}{2}$ ~ $\frac{2}{3}$ strip to beehive with 4~8 comb frames; Apply $\frac{2}{3}$ ~1 strip to beehive with more than 8 comb frames. Apply the strips one more time after 5~7 days. Applying medicine 3~4 times is a course of treatment

TRIAL IS REQUIRED

Choose 3 or 5 bee colonies for trial before applying to the whole colonies. Make a specific medicine using plan after observing 24~48 hours of the trial. It's better to conduct the trial in the morning of sunny day and adjust dose according to actual situation. Increase or reduce dose or medicine applying time on the condition that there are step hives, less comb frames or severe varroa mites disease



STEP 1



STEP 2

 Precaution

1. Please air-dry the strips in the shade for 24~48 hours if you find the strips are wet or sticky to each other. Strips can be come into use when there's no stickiness to fingers, otherwise it may cause harm to bees. Please do not place it under the Sun, otherwise the efficacy will be greatly decreasing.
2. Do not use this product when temperature is above 35°C or below 15°C. The product is not recommended to be applied to the situation such as migratory beekeeping, the beginning of hiving bee colonies, sugar deficiency for bees or other abnormal situation .
3. The said product can not be applied to bee colonies that have been fed with alkaline pollen such as China camellia pollen or other alkaline medicines and products recently or in the period of camellia collection.
4. The strips should not be placed continuously in beehive for more than 6 weeks.
5. When one decides to hang strips, please wear protective gloves and mask. Wash hands after use.
6. Disused strips should not be reused and should be disposed in a centralized place.
7. The strips are prohibited from contact with edible honey and should not be stored together with pesticides or other chemicals.
8. Store the said product somewhere away from food and children's reach and only unpack the packet when it is near to use.
9. Do not use this product during honey flow period.